arXiv:1210.5044v1 [math.DG] 18 Oct 2012

COVARIANT ALMOST ANALYTIC VECTOR FIELD ON Q - QUASI UMBILICAL HYPERSURFACE OF A SASAKIAN MANIFOLD

By

Sachin Kumar Srivastava, Alok Kumar Srivastava and Dhruwa Narain

Abstract : In this paper we have studied the properties of covariant almost analytic vector field on Q - quasi umbilical hypersurface M of a Sasakian manifold $\tilde{M}$ with $(\phi, g, u, v, \lambda)-$structure and obtained the scalars α and β using $1-form u, v$ covariant almost analytic for the hypersurface M to be totally umbilical and cylindrical.

2000 Mathematics Subject Classification : 53D10, 53C25, 53C21.
Keywords : Covariant derivative, Hypersurface, Sasakian manifold.

1. Introduction : Let $\tilde{M}$ be a $(2n+1)-$dimensional Sasakian manifold with a tensor field ϕ of type (1, 1), a fundamental vector field ξ and $1-form\,\eta$ such that

(1.1) $\eta(\xi)=1$

(1.2) $\tilde{\phi}^2=-I+\eta\otimes\xi$

where I denotes the identity transformation.

(1.3)(a) $\eta\, o\, \phi=0$ (b) $\phi\xi=0$ (c) $rank(\tilde{\phi})=2n$

If $\tilde{M}$ admits a Riemannian metric $\tilde{g}$, such that

(1.4) $\tilde{g}(\tilde{\phi}X,\tilde{\phi}Y)=\tilde{g}(X,Y)-\eta(X)\eta(Y)$

(1.5) $\tilde{g}(X,\xi)=\eta(X)$

then $\tilde{M}$ is said to admit a $(\tilde{\phi},\xi,\eta,\tilde{g})-$ structure called contact metric strucure.

If moreover,

(1.6) $(\tilde{\nabla}_X\tilde{\phi})\,Y=\tilde{g}(X,\,Y)\xi-\eta(Y)X$

and

(1.7) $\tilde{\nabla}_X\xi=-\tilde{\phi}X$

where $\tilde{\nabla}$ denotes the Riemannian connection of the Riemannian metric g, then $(\tilde{M},\tilde{\phi},\xi,\eta,\tilde{g})$ is called a Sasakian manifold [9]. If we define $'F(X,Y)=g(\phi X,Y)$, then in addition to above relation we find

(1.8) $'F(X,Y)+{}'F(Y,X)=0$

(1.9) $'F(X,\phi Y)={}'F(Y,\phi X)$

(1.10) $'F(\phi X,\phi Y)={}'F(X,Y)$

2. Hypersurface of a Sasakian manifold with $(\phi,g,u,v,\lambda)-$structure :

Let us consider a $2n-$dimensional manifold M embedded in $\tilde{M}$ with embedding $b: M\to\tilde{M}$. The map b induces a linear transformation map B (called Jacobian map), $B: T_p\to T_{b_p}$. Let an affine normal N of M is in such a way that $\tilde{\phi}N$ is always tangent to the hypersurface and satisfying the linear transformations

(2.1) $\tilde{\phi}BX=B\phi X+u(X)N$

(2.2) $$\tilde{\phi}N = -BU$$
(2.3) $$\xi = BV + \lambda N$$
(2.4) $$\eta(BX) = v(X)$$

where ϕ is a (1, 1) type tensor; U, V are vector fields; u, v are $1-form$ and λ is a $C^\infty-$ function. If $u \neq 0$, M, is called a noninvariant hypersurface of $\tilde{M}$ [1].

Operating (2.1), (2.2), (2.3) and (2.4) by $\tilde{\phi}$ and using (1.1), (1.2) and (1.3) and taking tangent normal parts separately, we get the following induced structure on M,

(2.5)(a) $$\phi^2 X = -X + u(X)U + v(X)V$$

(b) $$u(\phi X) = \lambda v(X), \qquad v(\phi X) = -\,\eta(N)\,u(X)$$

(c) $$\phi U = -\,\eta(N)\,V, \qquad \phi V = \lambda U$$

(d) $$u(U) = 1 - \lambda\eta(N), \qquad u(V) = 0$$

(e) $$v(U) = 0, \qquad v(V) = 1 - \lambda\eta(N)$$

and from (1.4) and (1.5), we get the induced metric g on M, i.e.,

(2.6) $$g(\phi X, \phi Y) = g(X, Y) - u(X)u(Y) - v(X)v(Y)$$
(2.7) $$g(U, X) = u(X), \qquad g(V, X) = v(X).$$

If we consider $\eta(N) = \lambda$, we get the following structures on M.

(2.8)(a) $$\phi^2 = -I + u \otimes U + v \otimes V$$

(b) $$\phi U = -\,\lambda V, \qquad \phi V = \lambda U$$

(c) $$u \circ \phi = \lambda v, \qquad v \circ \phi = -\lambda u$$

(d) $$u(U) = 1 - \lambda^2, \qquad u(V) = 0$$

(e) $$v(U) = 0, \qquad v(V) = 1 - \lambda^2.$$

A manifold M with a metric g satisfying (2.6), (2.7) and (2.8) is called manifold with $(\phi, g, u, v, \lambda)-$structure[2]. Let ∇ be the induced connection on the hypersurface M of the affine connection $\tilde{\nabla}$ of $\tilde{M}$.

Now using Gauss and Weingarten's equations

(2.9) $$\tilde{\nabla}_{BX} BY = B\nabla_X Y + h(X, Y)N$$

(2.10) $$\tilde{\nabla}_{BX} N = BHX + w(X)N, \text{ where } g(HY, Z) = h(Y, Z).$$

Here h and H are the second fundamental tensors of type (0, 2) and (1, 1) and w is a $1-form$. Now differentiating (2.1), (2.2), (2.3) and (2.4) covariantly and using (2.9), (2.10), (1.6) and reusing (2.1), (2.2), (2.3) and (2.4), we get

(2.11) $$(\nabla_Y\phi)(X) = v(X)Y - g(X,Y)V - h(X,Y)U - u(X)HY$$

(2.12) $$(\nabla_Y u)(X) = -h(\phi X, Y) - u(X)w(Y) - \lambda g(X,Y)$$

(2.13) $$(\nabla_Y v)(X) = g(\phi Y, X) + \lambda h(X,Y)$$

(2.14) $$\nabla_Y U = w(Y)U - \phi HY - \lambda Y$$

(2.15) $$\nabla_Y V = \phi Y + \lambda HY$$

(2.16) $$h(Y,V) = u(Y) - Y\lambda - \lambda w(Y)$$

(2.17) $$h(Y,U) = -u(HY)$$

Since $h(X,Y) = g(HX,Y)$, then from (1.5), and (2.17), we get

(2.18) $$h(Y,U) = 0 \Rightarrow HU = 0.$$

3. Q - Quasi Umbilical hypersurface : If

(3.1) $$h(X,Y) = \alpha g(X,Y) + \beta q(X)q(Y)$$

where α, β are scalar functions, q is $1-form$, then M is called Quasi-umbilical hypersurface and if $g(Q,X) = q(X)$, where Q is vector field, then M is called $Q-$quasi umbilical hypersurface. If $\alpha = 0$, $\beta \neq 0$, then $Q-$quasi umbilical hypersurface M is called cylindrical hypersurface. If $\alpha \neq 0$, $\beta = 0$, then $Q-$quasi-umbilical hypersurface M is called totally umbilical and if $\alpha = 0, \beta = 0$, then $Q-$quasi umbilical hypersurface is totally geodesic [9].

Using (3.1) in (2.11), (2.12), (2.13), (2.14), (2.15), (2.16) and (2.17) we get

(3.2) $$(\nabla_Y\phi)(X) = v(X)Y - g(X,Y)V - \{\alpha g(X,Y) + \beta q(X)q(Y)\}U$$
$$-u(X)\{\alpha Y + \beta q(Y)Q\}$$

(3.3) $$(\nabla_Y u)(X) = -\{\alpha g(\phi X, Y) + \beta q(\phi X)q(Y)\} - u(X)w(Y) - \lambda g(X,Y)$$

(3.4) $$(\nabla_Y v)(X) = g(\phi Y, X) + \lambda\{\alpha g(X,Y) + \beta q(X)q(Y)\}$$

(3.5) $$\nabla_Y U = w(Y)U - \{\alpha\phi Y + \beta q(Y)Q\} - \lambda Y$$

(3.6) $$\nabla_Y V = \phi Y + \lambda\{\alpha Y + \beta q(Y)Q\}$$

(3.7) $$h(Y,V) = \alpha g(V,Y) + \beta q(V)q(Y)$$

(3.8) $$|u(Q)|^2 = -\frac{\alpha}{\beta}(1-\lambda^2)$$

Also from (3.1), (2.16) and (2.8)(d) we get

(3.9). $$w(U) = \frac{1-\lambda^2}{\lambda} - \frac{U\lambda}{\lambda}$$

4. Covariant almost analytic vector field on $Q-$quasi umbilical hypersurface : $1-form$ u and v are said to be covariant almost analytic if

$$u\{(\nabla_X\phi)(Y)-(\nabla_Y\phi)(X)\}=(\nabla_{\phi X}u)(Y)-(\nabla_X u)(\phi Y) \tag{4.1}$$

and

$$v\{(\nabla_X\phi)(Y)-(\nabla_Y\phi)(X)\}=(\nabla_{\phi X}v)(Y)-(\nabla_X v)(\phi Y) \tag{4.2}$$

Theorem 4.1 : *On $Q-$quasi umbilical hypersurface M with $(\phi,g,u,v,\lambda)-$ structure of a Sasakian manifold $\tilde{M}$, if $1-form$ u is covariant almost analytic, then we have*

$$\begin{aligned} v(Y)u(X)-v(X)u(Y) = &-2\alpha g(\phi X,\phi Y)-2\lambda g(\phi X,Y)-\beta q(\phi Y)q(\phi X)\\ &-\beta q(Y)q(X)+2\beta q(X)u(Y)u(Q)\\ &-\beta u(X)q(Y)-u(Y)w(\phi X)+u(\phi Y)w(X)\end{aligned} \tag{4.3}$$

Proof : From (3.2), we have

$$\begin{aligned}(\nabla_Y\phi)(X) = v(X)Y-g(X,Y)V-\{\alpha g(X,Y)+\beta q(X)q(Y)\}U\\ -u(X)\{\alpha Y+\beta q(Y)Q\}\end{aligned}$$

and
$$\begin{aligned}(\nabla_X\phi)(Y) = v(Y)X-g(Y,X)V-\{\alpha g(Y,X)+\beta q(Y)q(X)\}U\\ -u(Y)\{\alpha X+\beta q(X)Q\}\end{aligned}$$

$$\therefore \quad \begin{aligned}(\nabla_X\phi)(Y)-(\nabla_Y\phi)(X) = v(Y)X-v(X)Y-u(Y)\{\alpha X+\beta q(X)Q\}\\ +u(X)\{\alpha Y+\beta q(Y)Q\}\end{aligned}$$

$$\begin{aligned}u\{(\nabla_X\phi)(Y)-(\nabla_Y\phi)(X)\} = v(Y)u(X)-v(X)u(Y)\\ +\beta\{u(X)q(Y)-q(X)u(Y)\}u(Q)\end{aligned} \tag{4.4}$$

Also from (3.4), we have

$$(\nabla_X u)(Y)=-\{\alpha g(\phi Y,X)+\beta q(\phi Y)q(X)\}-u(Y)w(X)-\lambda g(X,Y)$$

or
$$(\nabla_{\phi X}u)(Y)=-\alpha g(\phi X,\phi Y)-\beta q(\phi Y)q(\phi X)-u(Y)w(\phi X)-\lambda g(\phi X,Y)$$

and
$$(\nabla_X u)(\phi Y)=-\alpha g(\phi^2 Y,X)-\beta q(\phi^2 Y)q(X)-u(\phi Y)w(X)-\lambda g(X,\phi Y)$$

or
$$\begin{aligned}(\nabla_X u)(\phi Y)=\alpha g(Y,X)-\alpha u(X)u(Y)-\alpha v(X)v(Y)+\beta q(Y)q(X)\\ -\beta u(Y)q(U)q(X)-\beta v(Y)q(V)q(X)-u(\phi Y)w(X)-\lambda g(X,\phi Y)\end{aligned}$$

or
$$\begin{aligned}(\nabla_X u)(\phi Y)=\alpha g(\phi X,\phi Y)+\beta q(Y)q(X)-\beta u(Y)q(U)q(X)\\ -u(\phi Y)w(X)-\lambda g(X,\phi Y)\end{aligned}$$

$$\begin{aligned}(\nabla_{\phi X}u)(Y)-(\nabla_X u)(\phi Y) = &-2\alpha g(\phi X,\phi Y)-\beta q(\phi Y)q(\phi X)-\beta q(Y)q(X)\\ &+\beta u(Y)q(U)q(X)-u(Y)w(\phi X)+u(\phi Y)w(X)\\ &-\lambda g(\phi X,Y)+\lambda g(X,\phi Y)\end{aligned} \tag{4.5}$$

using (4.1) in (4.4) and (4.5), we get (4.3).

Corollary 4.1 : *On $Q-$quasi umbilical hypersurface M with $(\phi,g,u,v,\lambda)-$ structure of a Sasakian manifold $\tilde{M}$, if $1-form$ u is covariant almost analytic, then we also* *have*

(4.6) $$\nabla_U \lambda = 2\lambda^2.$$

Proof : Put $X = U$ and $Y = V$ in (4.3) and using (2.8)(d), (2.8)(e), we get

$$(1-\lambda^2)^2 \;=\; -2\,\alpha.0 + 2\lambda^2(1-\lambda^2) - \beta.0 - 0 + \lambda(1-\lambda^2)\,w(U).$$

Using (3.9), we get

$$(1-\lambda^2) = 2\lambda^2 + \lambda\left\{\frac{1-\lambda^2}{\lambda} - \frac{(U\lambda)}{\lambda}\right\}$$

or $$(1-\lambda^2) - 2\,\lambda^2 - 1 + \lambda^2 = -\,U\lambda$$

or $$2\,\lambda^2 = U\lambda$$

or $$U\lambda = 2\,\lambda^2$$

or $$\nabla_U\lambda = 2\lambda^2.$$

Theorem 4.2 : *On cylindrical hypersurface M with* $(\phi, g, u, v, \lambda)-$*structure of a Sasakian manifold* $\tilde{M}$*, with covariant almost analytic vector fields* U*,* V*, we have*

(4.7) $$\begin{aligned} v(Y)u(X) - v(X)u(Y) = {}& -2\lambda g(\phi X, Y) - \beta q(\phi X)q(\phi Y) - \beta q(Y)q(X) \\ & - \beta u(X)q(Y) - u(Y)w(\phi X) + u(\phi Y)w(X) \end{aligned}$$

Proof : Putting $\alpha = 0$ in (4.3) and using $u(Q) = 0$, we get (4.7).

Corollary 4.2 : *On cylindrical hypersurface M with* $(\phi, g, u, v, \lambda)-$*structure of a Sasakian manifold* $\tilde{M}$*, with covariant almost analytic vector fields* U*,* V*, we also have*

(4.8) $V\lambda = 0$, i.e. λ is covariant constant along V

(4.9) $$q(Q) = 0.$$

Proof : Putting $X = V$ in (4.7), we get

$$\begin{aligned} -\,(1-\lambda^2)u(Y) = {}& -\,2\lambda g(\phi V, Y) - \beta q(\phi Y)q(\phi V) - \beta q(V)q(Y) \\ & - \beta u(V)q(Y) - u(Y)w(\phi V) + u(\phi Y)w(V) \end{aligned}$$

or $$-\,(1-\lambda^2)u(Y) = -\,2\lambda^2 u(Y) - \lambda u(Y)w(U) + u(\phi Y)w(V)$$

or $$(3\lambda^2 - 1)u(Y) = -\,\lambda u(Y)\left(\frac{1-\lambda^2}{\lambda} - \frac{U\lambda}{\lambda}\right) + u(\phi Y)\left(\frac{-\,V\lambda}{\lambda}\right)$$

or $$(3\lambda^2 - 1)u(Y) = -\,u(Y)\{(1-\lambda^2) - (U\lambda)\} - u(\phi Y)\left(\frac{V\lambda}{\lambda}\right)$$

or $$(3\lambda^2 - 1 + 1 - \lambda^2)U = (U\lambda)U + \left(\frac{V\lambda}{\lambda}\right)\phi U$$

or $$2\lambda^2\, U = (U\lambda)\, U + \left(\frac{V\,\lambda}{\lambda}\right)\phi U$$

using (4.6), we get $V\lambda = 0$, i.e. λ is covariant constant along V.

Also from (4.7), we get

$$\begin{aligned} u(X)V - v(X)U = &-2\lambda\phi X + \beta\phi Q q(\phi X) - \beta q(X)Q - \beta u(X)Q \\ &- w(\phi X)\, U - w\,(X)\phi U \end{aligned}$$

Contracting with respect to X, we get

$$0 = \beta q(\phi^2 Q) - \beta q(Q) - \beta u(Q) + \lambda w(V) + \lambda w(V)$$

or $$\beta q(Q) = -(V\lambda) = 0$$

since $\beta \neq 0$, so $q(Q) = 0$.

Theorem 4.3 : *On $Q-$quasi umbilical hypersurface M with $(\phi, g, u, v, \lambda)-$structure of a Sasakian manifold $\tilde{M}$, where $1-formu$ is covariant almost analytic. If M is totally umbilical, then we have*

$$\begin{aligned} v(Y)u(X) - v(X)u(Y) = &-2\alpha g(\phi X, \phi Y) - 2\lambda g(\phi X, Y) \\ &- u(Y)w(\phi X) + u(\phi Y)w(X) \end{aligned} \tag{4.10}$$

$$\alpha = -\tfrac{(V\lambda)}{1+\lambda^2} \quad \text{or} \quad \alpha = -\,d(\tan^{-1}\lambda)\, V. \tag{4.11}$$

Proof : Putting $\beta = 0$ in (4.3), we get (4.10). Further putting $X = U$ in (4.10), we get

$$(1-\lambda^2)v(Y) = 2\,\alpha\lambda v(\phi Y) + 2\lambda^2 v(Y) + \lambda u(Y)w(V) + u(\phi Y)w(U)$$

$$\begin{aligned} (1-\lambda^2)v(Y) = &-2\,\alpha\lambda^2 u(Y) + 2\lambda^2 v(Y) \\ &+ \lambda u(Y)\left\{\alpha\frac{(\lambda^2-1)}{\lambda} - \frac{(V\lambda)}{\lambda}\right\} + u(\phi Y)\left\{\frac{1-\lambda^2}{\lambda} - \frac{(U\lambda)}{\lambda}\right\} \end{aligned}$$

or $$\begin{aligned} (1-\lambda^2-2\lambda^2-1+\lambda^2)v(Y) = &\{-2\alpha\lambda^2 + \alpha\lambda^2 - \alpha\}\, u(Y) - u(Y)(V\lambda) \\ &-v(Y)(U\lambda) \end{aligned}$$

or $$-2\lambda^2 V = -\,\alpha\,(1+\lambda^2)U - (V\lambda)\, U - (U\,\lambda)\, V$$

Operating by u

or $$0 = -\,\alpha\,(1+\lambda^2)\,(1-\lambda^2) - (V\lambda)\,(1-\lambda^2)$$

or $$\alpha = -\frac{(V\,\lambda)}{1+\lambda^2}$$

or $$\alpha = -\,d\,(\tan^{-1}\lambda)\,(V)$$

Corollary 4.3 : *On Q−quasi umbilical hypersurface M with* (ϕ, g, u, v, λ)−*structure of a Sasakian manifold* $\tilde{M}$, *where* $1-form$ *u is covariant almost analytic. If M is totally umbilical, then we also have*

$$\alpha = -\frac{1}{\sqrt{2n-1}} d\left(\tan^{-1}\frac{\lambda}{\sqrt{2n-1}}\right)(V) \tag{4.12}$$

Proof : From (4.10), we have

$$u(X)V - v(X)U = 2\alpha\phi^2X - 2\lambda\phi X - w(\phi X)U + \lambda w(X)V$$

Contracting with respect to X

$$0 = 2\,\alpha\{-2n + 4\alpha(1-\lambda^2)\} + 2\lambda w(V)$$

or $\quad 0 = -2\,\alpha\,n + 2\alpha(1-\lambda^2) + \lambda w(V)$

or $\quad -2\,\alpha\,n + 2\alpha(1-\lambda^2) + \lambda\left\{\alpha\frac{(\lambda^2-1)}{\lambda} - \frac{(V\lambda)}{\lambda}\right\} = 0$

or $\quad \alpha(1-\lambda^2-2n) = (V\lambda)$

or $\quad \alpha = \dfrac{V\lambda}{1-\lambda^2-2n} = -\dfrac{(d\lambda)V}{(2n-1)+\lambda^2}$

thus $\quad \alpha = -\dfrac{1}{\sqrt{2n-1}} d\left(\tan^{-1}\dfrac{\lambda}{\sqrt{2n-1}}\right)(V).$

Theorem 4.4 : *On a Q−quasi umbilical noninvariant hypersurface M with* (ϕ, g, u, v, λ)−*structure of a Sasakian manifold* $\tilde{M}$, *If* $1-form$ *v is covariant almost analytic, then we have*

$$\begin{aligned}\alpha\{u(X)v(Y) - u(Y)v(X)\} = &-2g(\phi X, \phi Y) + 2\lambda\alpha g(\phi X, Y)\\ &+ \lambda\beta\{q(\phi X)q(Y) - q(X)q(\phi Y)\}\end{aligned} \tag{4.13}$$

Proof : From (3.2), we get

$$\begin{aligned}(\nabla_X\phi)(Y) - (\nabla_Y\phi)(X) = v(Y)X - v(X)Y - &u(Y)\{\alpha X + \beta q(X)Q\}\\ &+ u(X)\{\alpha Y + \beta q(Y)Q\}\end{aligned}$$

$$v\{(\nabla_X\phi)(Y) - (\nabla_Y\phi)(X)\} = \alpha\{u(X)\,v(Y) - u(Y)\,v(X)\} \tag{4.14}$$

from (3.4), we get

$$(\nabla_{\phi X}v)(Y) = -g(X, Y) + u(X)u(Y) + v(X)v(Y) + \lambda\alpha g(\phi X, Y) + \lambda\beta q(\phi X)q(Y)$$

and

$$(\nabla_X v)(\phi Y) = g(X, Y) - u(X)u(Y) - v(X)v(Y) + 2\lambda g(X, \phi Y) + \lambda\beta q(X)q(\phi Y)$$

therefore

$$(4.15)\quad (\nabla_{\phi X} v)(Y) - (\nabla_X v)(\phi Y) = -\,2\,g(\phi X, \phi Y) + \lambda\beta\{q(\phi X)q(Y) - q(X)q(\phi Y)\} \\ +2\lambda\alpha g(\phi X, Y)$$

Using (4.2) in (4.14) and (4.15), we get (4.13).

Theorem 4.5 : *Let $1-form$ v is covariant almost analytic on $Q-$quasi umbilical noninvariant hypersurface M with $(\phi, g, u, v, \lambda)-$structure of a Sasakian manifold $\tilde{M}$, and if it is cylindrical also, we have*

$$(4.16)\qquad 2g(\phi X, \phi Y) = \lambda\beta\{q(\phi X)q(Y) - q(X)q(\phi Y)\}, \text{ with } X \neq U.$$

Proof : Putting $\alpha = 0$ in (4.13), we get (4.16). Further if $X = U$, then from (4.16), we get

$$2g(\phi U, \phi Y) = \lambda\beta\{-\,\lambda q(V)q(Y) - q(U)q(\phi Y)\}$$

or $$-\,2\lambda g\,(V, \phi Y) = 0$$

or $$-\,2\lambda v\,(\phi Y) = 0$$

since $v(\phi Y) \neq 0$ and $\lambda \neq 0$, therefore $X \neq U$.

Theorem 4.6 : *Let $1-form$ v is covariant almost analytic on $Q-$quasi umbilical noninvariant hypersurface M with $(\phi, g, u, v, \lambda)-$structure of a Sasakian manifold, and if it is totally umbilical, then we have*

(4.17) $\alpha\{u(X)v(Y) - v(X)u(Y)\} = -2\,g(\phi X, \phi Y) + 2\lambda\alpha g(\phi X, Y)$, with $X \neq U$.

Proof : Putting $\beta = 0$ in (3.13), we get (3.17). If we take $X = U$ in (4.17), we get

$$\alpha\,(1 - \lambda^2)\,v(Y) = 2\,\lambda v(\phi Y) - 2\,\lambda\alpha v(Y)$$

or $$\alpha\,\{1 - \lambda^2 + 2\lambda\}\,v(Y) = 2\,\lambda v(\phi Y)$$

or $$\alpha\,\{\lambda^2 - 2\lambda - 1\}\,v(Y) = -\,2\,\lambda\phi V = -\,2\lambda^2 U$$

i.e. V and U are linearly dependent, which is a contradiction. Thus $X \neq U$.

References

[1] S. I. Goldberg and K. Yano : *Noninvariant hypersurfaces of almost contact manifolds*, J. Math. Soc., Japan, **22** (1970), 25-34.

[2] D. Narain : *Hypersurfaces with $(f, g, u, v, \lambda)-$structure of an affinely cosymplectic manifold*, Indian Jour. Pure and Appl. Math.**20** No. 8 (1989), 799-803.

[3] Dhruwa Narain and S. K. Srivastva : *Noninvariant Hypersurfaces of Sasakian Space Forms*, Int. J. Contemp. Math. Sciences, Vol. **4**, 2009, no. 33, 1611 - 1617.

[4] Dhruwa Narain, S. K. Srivastava and Khushbu Srivastava : *A note of noninvariant hypersurfaces of para Sasakian manifold*, IOSR Journal of Engineering(IOSRJEN) Vol.**2**, issue 2 (2012), 363-368.

[5] Dhruwa Narain and S. K. Srivastava : *On the hypersurfaces of almost r-contact manifold*, J. T. S. Vol.**2**(2008), 67-74.

[6] Rajendra Prasad and M. M. Tripathi : *Transversal Hypersurfaces of Kenmotsu Manifold*, Indian Jour. Pure and Appl. Math.**34** No. 3 (2003), 443-452.

[7] B. B. Sinha and Dhruwa Narain : *Hypersurfaces of nearly Sasakian manifold*, Ann. Fac. Sci. Dekishasa Zaire, **3** (2), (1977), 267-280.

[8] M. Okumura : *Totally umbilical hypersurface of a locally product Riemannian manifold*, Kodai Math. Sem. Rep.**19** (1967), 35-42.

[9] D. E. Blair : *Almost contact manifolds with killing structure tensor*, Pacific J. of Math. **39** (12), (1971), 285-292.

[10] T. Miyazawa and S. Yamaguchi : *Some theorems on K-contact metric manifolds and Sasakian manifolds*, T. R. V. Math. **2** (1966), 40-52.

Authors' addresses:
Sachin Kumar Srivastava
Amity Institute of Applied Sciences,Amity University , Noida, U.P., India
$E - mail$: *sksrivastava4@amity.edu*

Alok Kumar Srivastava
Department of Mathematics ,Govt. Degree College, Chunar - Mirzapur,U.P., India
$E - mail$: *aalok_sri@yahoo.co.in*

Dhruwa Narain
Department of Mathematics and Statistics
D.D.U. Gorakhpur University Gorakhpur, Gorakhpur-273009, India
$E - mail$: *dhruwanarain_dubey@yahoo.co.in*